






























Dosage Chart

VETMEDIN should be administered orally at a total daily dose of 0.5mg/kg body weight, divided into two portions (that are not necessarily equal) given approximately 12 hours apart, using a suitable combination of whole or half tablets.

NUMBER OF CHEWABLE TABLETS
PER ADMINISTRATION

		WEIGHT	TOTAL DAILY DOSAGE	MORNING 	EVENING 
1.25mg		2.5kg	1.25mg		
		3.75kg	1.875mg		
		5kg	2.5mg		
		7.5kg	3.75mg		
5.0mg		10kg	5mg		
		15kg	7.5mg		
		20kg	10mg		
		30kg	15mg		
		40kg	20mg		

VETMEDIN should be administered orally at a total daily dose of 0.5mg/kg body weight, using a suitable combination of whole or half tablets. The total daily dose should be divided into two portions that are not necessarily equal, and the portions should be administered approximately 12 hours apart (i.e., morning and evening). The tablets are scored and the calculated dosage should be provided to the nearest half tablet increment.

The examples given in the dosage chart do not necessarily depict the only way to provide the correct dose. Other combinations of whole and half tablets may also be appropriate.

Please refer to the full prescribing information on the back of this chart. For complete product information visit www.vetmedin-us.com.

Important Safety Information: VETMEDIN should not be given in case of hypertrophic cardiomyopathy, aortic stenosis or any other clinical condition where an augmentation of cardiac output is inappropriate for functional or anatomical reasons. The safety of VETMEDIN has not been established in dogs with asymptomatic heart disease or in heart failure caused by other etiologies other than atrioventricular valvular insufficiency or dilated cardiomyopathy. Use only in dogs with clinical evidence of heart failure. The most common side effects reported in field studies were poor appetite, lethargy, diarrhea, dyspnea, azotemia, weakness and ataxia. If side effects should occur, pet owners should contact their veterinarian.